

Product Information Sheet

MYCOVAC-L® ***Mycoplasma gallisepticum* Vaccine** **Live Culture**

Product Information

MYCOVAC-L® is the patented, live Intervet 6/85 strain of *Mycoplasma gallisepticum* (MG). The safe, highly immunogenic, lyophilized vaccine is for use via fine spray in chickens for protection against losses associated with MG infection.




MYCOVAC-L® contains:

- Patented modified live 6/85 strain of *Mycoplasma gallisepticum*.

Product Features:

- Minimal lesions, less stress, safer for chickens than F-strain vaccines.
- Not pathogenic to turkeys.
- Does not spread to adjacent chicken houses.
- No ovarian transmission.
- Protects against the highly virulent strains of MG.
- Protects against losses in egg production associated with MG.
- Serum will not convert MG plate antigen: easy to distinguish vaccine response from field infection.



Indications for Use:

Mycovac-L[®] is indicated for the vaccination of chickens at six weeks of age or older for protection against clinical signs of MG.

Dosage and Administration:

The time of administration should be determined by the degree of MG exposure, type of chicken being vaccinated, and the current management procedures. **Mycovac-L** is approved for use via fine spray (less than 20 microns) at 6 weeks of age or older.

Vaccination:

The best protection is obtained when the vaccinated chickens have time to develop immunity prior to exposure to MG. Avoid the use of any medications effective against MG for 5 days before or after vaccination. Good management practices are recommended to reduce the exposure to MG for at least 3 weeks following vaccination. Birds should be maintained in a proper environment with good management at all times to maximum vaccine efficacy.

Cautions:

1. Refer to the package insert for specific directions.
2. Do not medicate the chickens with antibacterial drugs--especially chlortetracycline, xytetracycline, and sulfonamid--5 days before or after vaccination.
3. This vaccine should not be administered within 2 weeks of any Newcastle, bronchitis or laryngotracheitis vaccination.
4. All susceptible birds on the same premises should be vaccinated at the same time.
5. Do not dilute the vaccine or otherwise stretch the dosage of the vaccine.
6. Use entire contents of the vial when first opened.
7. Do not vaccinate chickens within 21 days of slaughter.
8. The 6/85 strain is not pathogenic for turkeys, but care should be taken to maintain tight biosecurity during the application of the vaccine and when handling vaccinated chickens.

How Supplied

10 x 2,000 dose vials

Product Code

MG-240-2

Cap Color Code

